arXiv:quant-ph/9705007v2 2 Jul 1997

GTP-97-01

Propagator for an Aharonov-Bohm-Coulomb system

[a]D.K.Park, [b]Sahng-Kyoon Yoo, [c]Soo-Young Lee, [c]Jae-Rok Kahng,
[d]Chang Soo Park, [e]Eui-Soon Yim, and [f]C.H.Lee

[a] *Department of Physics, Kyungnam University, Masan, 631-701, Korea*

[b] *Department of Physics, Seonam University, Namwon, 590-170, Korea*

[c] *Department of Physics, Korea University, Seoul, 136-701, Korea*

[d] *Department of Physics, Dankook University, Cheonan, 330-180, Korea*

[e] *Department of Physics, Semyung University, Chechon, 390-230, Korea*

[f] *D&S Dept.,R&D Center, Anam Industrial CO., LTD, Seoul 133-120, Korea*

Abstract

The propagator of three-dimensional Aharonov-Bohm-Coulomb system is calculated by following the Duru-Kleinert method. It is shown that the system is reduced to two independent two dimensional Aharonov-Bohm plus harmonic oscillator systems through dimensional extension and Kustaanheimo-Stiefel transformation. The energy spectrum is deduced.

Since Aharonov and Bohm(AB) gave a physical significance to the vector potential in 1959[1], there are lots of theoretical and experimental attempts[2] to establish the AB effect. Recent attention given to the phenomenon is mostly related to the anyonic theory[3] which tries to understand the high-T_c superconductivity phenomena using fractional statistics. Since AB interaction is interaction between charge and magnetic flux, anyon-anyon interaction naturally requires Coulomb modification. In this letter we will study on the AB problem with Coulomb modification by invoking the path-integral formalism in three dimensions.

For a last decade, the two-[4] and three-dimensional[5,6] Aharonov-Bohm-Coulomb(ABC) systems have been studied by using the various different methods. Especially Ref.[5] used the path-integral formalism in the spherical coordinate and obtained the energy-dependent Green's function and the bound-state energy spectrum in ABC system. Same bound-state energy spectrum has been found in Ref.[6], in which the generalized ABC system is studied with an aid of Kustaanheimo-Stiefel(KS) transformation[7] by solving Schrödinger equation. The KS transformation is also used for the evaluation of the exact propagator in hydrogen atom[8,9].

In this letter we will derive the exact propagator for ABC system by invoking the Duru-Kleinert method. Let us start with Hamiltonian

$$\hat{H} = \frac{(\vec{p} - e\vec{A})^2}{2M} + \frac{\xi}{r} \tag{1}$$

where AB potential is

$$\vec{A} = \frac{\alpha}{e} \frac{y\hat{x} - x\hat{y}}{r^2 - z^2} \tag{2}$$

in Coulomb gauge. Following Ref.[10], we can describe the fixed-energy amplitude $(\vec{x}_b \mid \vec{x}_a)_E$ as a pseudotime-sliced path integral

$$\begin{aligned} &(\vec{x}_b \mid \vec{x}_a)_E \qquad\qquad (3) \\ &= \int_0^\infty ds f_r(\vec{x}_b) f_l(\vec{x}_a) \int D\vec{x} \int D\vec{p} exp \left[i \int_0^s ds \ \left[\vec{p} \cdot \vec{x}' - f_l(\vec{x})(\hat{H} - E) f_r(\vec{x}) \right] \right] \end{aligned}$$

where $f_l(\vec{x})$ and $f_r(\vec{x})$ are regulating functions defined in chapter 12 of Ref.[10] and $\vec{x}' = d\vec{x}/ds$. Pseudotime s is defined by $ds/dt = f_l(\vec{x})f_r(\vec{x})$. It is worthwhile to note that fixed-energy amplitude has a relation with energy-dependent Green's function , which is Laplace transform of Euclidean Kernel, as follows:

$$(\vec{x}_b \mid \vec{x}_a)_E = -i\hat{G}[\vec{x}_b, \vec{x}_a : -E]. \tag{4}$$

After choosing

$$\begin{aligned} f_l(\vec{x}) &= r^{1-\lambda} \\ f_r(\vec{x}) &= r^{\lambda} \end{aligned} \tag{5}$$

and performing $\vec{p}$-integration, one can derive the following form straightforwardly:

$$\begin{aligned} &(\vec{x}_b \mid \vec{x}_a)_E \\ &= \lim_{N\to\infty} (N+1) \int_0^\infty d\epsilon_s r_b^{\lambda} r_a^{1-\lambda} \left(\frac{M}{2\pi i\epsilon_s r_a^{\lambda} r_b^{1-\lambda}}\right)^{3/2} \int \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i\epsilon_s r_{j-1}}\right)^{3/2} d\Delta\vec{x}_j\right] e^{iA_{0,E}^N} \end{aligned} \tag{6}$$

where $(N+1)\epsilon_s = s_b - s_a \equiv s$ and

$$A_{0,E}^N = -\xi(N+1)\epsilon_s + \sum_{j=1}^{N+1} \left[\frac{M}{2} \frac{(\vec{x}_j - \vec{x}_{j-1})^2}{\epsilon_s r_j^{1-\lambda} r_{j-1}^{\lambda}} + e\vec{A}_j \cdot (\vec{x}_j - \vec{x}_{j-1}) + \epsilon_s E r_j\right] \tag{7}$$

whose contiuum limit is

$$A_{0,E}[\vec{x}] = -\xi s + \int_0^s ds \left[\frac{M}{2r}\vec{x}'^2 + e\vec{A}\cdot\vec{x}' + Er\right]. \tag{8}$$

At this stage we have to exploit the dimensional extension technique[11] for the incorporation of the KS transformation into the path integral formalism. This is achieved by inserting the following trivial identity

$$\int_{-\infty}^{\infty} \left[\prod_{j=1}^{N+1} \left(\frac{M}{2\pi i\epsilon_s r_j^{1-\lambda} r_{j-1}^{\lambda}}\right)^{1/2} d\Delta\eta_j\right] exp\left[i\sum_{j=1}^{N+1} \frac{M}{2} \frac{\Delta\eta_j^2}{\epsilon_s r_j^{1-\lambda} r_{j-1}^{\lambda}}\right] = 1 \tag{9}$$

into Eq.(6). Hence, $(\vec{x}_b \mid \vec{x}_a)_E$ is expressed as four dimensional path integral

$$\begin{aligned} (\vec{x}_b \mid \vec{x}_a)_E = (N+1) \int_0^\infty d\epsilon_s r_b^{\lambda} r_a^{1-\lambda} \left(\frac{M}{2\pi i\epsilon_s r_a^{\lambda} r_b^{1-\lambda}}\right)^2 \\ \times \int d\Delta\eta_1 \int \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i\epsilon_s r_{j-1}}\right)^2 d\Delta\vec{x}_j d\Delta\eta_j\right] e^{iA_E^N} \end{aligned} \tag{10}$$

where

$$A_E^N = -\xi(N+1)\epsilon_s + \sum_{j=1}^{N+1} \left[\frac{M}{2} \frac{\Delta\vec{x}_j^2 + \Delta\eta_j^2}{\epsilon_s r_j^{1-\lambda} r_{j-1}^{\lambda}} + e\vec{A}_j \cdot \Delta\vec{x}_j + \epsilon_s E r_j \right]. \tag{11}$$

By using the the following approximation

$$\begin{aligned} & r_b^{\lambda} r_a^{1-\lambda} \left(\frac{M}{2\pi i \epsilon_s r_a^{\lambda} r_b^{1-\lambda}} \right)^2 \prod_{j=2}^{N+1} \left(\frac{M}{2\pi i \epsilon_s r_{j-1}} \right)^2 \\ \approx & \left(\frac{M}{2\pi i \epsilon_s} \right)^2 \frac{1}{r_a} \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i \epsilon_s r_j} \right)^2 \right] exp \left[3\lambda \sum_{j=1}^{N+1} \ln \frac{r_j}{r_{j-1}} \right] \end{aligned} \tag{12}$$

one can change Eq.(10) into

$$\begin{aligned} (\vec{x}_b \mid \vec{x}_a)_E = & (N+1) \int_0^{\infty} d\epsilon_s \left(\frac{M}{2\pi i \epsilon_s} \right)^2 \\ & \times \int \frac{d\Delta\eta_1}{r_a} \int \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i \epsilon_s r_j} \right)^2 d\Delta\vec{x}_j d\Delta\eta_j \right] e^{i(A_E^N + A_f^N)} \end{aligned} \tag{13}$$

where

$$A_f^N = -3i\lambda \sum_{j=1}^{N+1} \ln \frac{r_j}{r_{j-1}}. \tag{14}$$

Since final continuum limit of $(\vec{x}_b \mid \vec{x}_a)_E$ is independent of λ [10], we set $\lambda = 0$, which gives

$$(\vec{x}_b \mid \vec{x}_a)_E = (N+1) \int_0^{\infty} d\epsilon_s \left(\frac{M}{2\pi i \epsilon_s} \right)^2 \int \frac{d\Delta\eta_1}{r_a} \int \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i \epsilon_s r_j} \right)^2 d\Delta\vec{x}_j d\Delta\eta_j \right] e^{iA^N_{E,\lambda=0}}. \tag{15}$$

We now apply the KS transformation defined by

$$\begin{pmatrix} \Delta x \\ \Delta y \\ \Delta z \\ \Delta\eta \end{pmatrix} = 2A(\vec{u}) \begin{pmatrix} \Delta u^1 \\ \Delta u^2 \\ \Delta u^3 \\ \Delta u^4 \end{pmatrix}, \tag{16}$$

where

$$A(\vec{u}) = \begin{pmatrix} u^3 & u^4 & u^1 & u^2 \\ u^4 & -u^3 & -u^2 & u^1 \\ u^1 & u^2 & -u^3 & -u^4 \\ u^2 & -u^1 & u^4 & -u^3 \end{pmatrix} \tag{17}$$

to the path integral calculation of $(\vec{x}_b \mid \vec{x}_a)_E$. ¿From the definition of KS transformation (16) and (17), it is easily shown that

$$\begin{aligned} x^2 + y^2 + x^2 &= \left[(u^1)^2 + (u^2)^2 + (u^3)^2 + (u^4)^2\right]^2, \\ \Delta x^2 + \Delta y^2 + \Delta z^2 + \Delta \eta^2 &= 4\left[(u^1)^2 + (u^2)^2 + (u^3)^2 + (u^4)^2\right] \\ &\quad \times \left[(\Delta u^1)^2 + (\Delta u^2)^2 + (\Delta u^3)^2 + (\Delta u^4)^2\right], \\ \frac{\partial(x, y, z, \eta)}{\partial(u^1, u^2, u^3, u^4)} &= 2^4 r^2. \end{aligned} \tag{18}$$

Furthermore, the KS transformation of AB potential term is

$$\begin{aligned} \vec{A} \cdot d\vec{x} &\equiv \frac{\alpha}{e} \frac{y dx - x dy}{x^2 + y^2} \\ &= \frac{\alpha}{e} \left[\frac{u^1 du^2 - u^2 du^1}{(u^1)^2 + (u^2)^2} + \frac{u^4 du^3 - u^3 du^4}{(u^3)^2 + (u^4)^2} \right], \end{aligned} \tag{19}$$

which makes the system separable like $R^4 \to R^2 \times R^2$. This separability from R^4 to two independent two-dimensional AB plus harmonic oscillator systems makes the path-integral calculation of $(\vec{x}_b \mid \vec{x}_a)_E$ extremely simple. Although $\vec{u}$-space is noneuclidean with curvature and torsion[12], H.Kleinert showed that one can change

$$\begin{aligned} &\int \frac{d\Delta\eta_1}{r_a} \int \left[\prod_{j=2}^{N+1} \left(\frac{M}{2\pi i \epsilon_s r_j} \right)^2 d\Delta\vec{x}_j d\Delta\eta_j \right] \\ &\qquad \Longrightarrow \int \frac{d\eta_a}{r_a} \int \left[\prod_{j=1}^{N} \left(\frac{4M}{2\pi i \epsilon_s} \right)^2 d^4\vec{u}_j \right] \end{aligned} \tag{20}$$

without time slicing correction. After inserting Eq.(20) into (15) and using Eqs.(16, 17, 18), one can perform the path integral which gives

$$\begin{aligned} (\vec{x}_b \mid \vec{x}_a)_E = 2^{-4} \int_0^\infty & ds e^{-i\xi s} \int \frac{d\eta_a}{r_a} \left(\frac{4M\omega}{2\pi i \sin\omega s} \right)^2 \\ &\times \sum_{m_1=-\infty}^{\infty} \sum_{m_2=-\infty}^{\infty} e^{im_1(\theta_{1,b} - \theta_{1,a})} e^{im_2(\theta_{2,b} - \theta_{2,a})} \\ &\times \exp\left[2iM\omega \frac{\cos\omega s}{\sin\omega s} (\rho_{1,a}^2 + \rho_{1,b}^2 + \rho_{2,a}^2 + \rho_{2,b}^2) \right] \\ &\times I_{|m_1+\alpha|} \left(\frac{-4iM\omega}{\sin\omega s} \rho_{1,a}\rho_{1,b} \right) I_{|m_2+\alpha|} \left(\frac{-4iM\omega}{\sin\omega s} \rho_{2,a}\rho_{2,b} \right) \end{aligned} \tag{21}$$

where (ρ_1, θ_1) and (ρ_2, θ_2) are double polar coordinates defined by

$$\begin{aligned} u^1 &= \rho_1 \sin\theta_1 \\ u^2 &= \rho_1 \cos\theta_1 \\ u^3 &= \rho_2 \cos\theta_2 \\ u^4 &= \rho_2 \sin\theta_2 \end{aligned} \tag{22}$$

and

$$\omega^2 = -\frac{E}{2M}. \tag{23}$$

In order to perform the η_a-integration we express $(\rho_1, \rho_2, \theta_1, \theta_2)$ in terms of three-dimensional spherical coordinate with an auxiliary angle γ:

$$\begin{aligned} \rho_1 &= \sqrt{r}\cos\frac{\theta}{2} \\ \theta_1 &= \frac{\phi+\gamma+\pi}{2} \\ \rho_2 &= \sqrt{r}\sin\frac{\theta}{2} \\ \theta_2 &= \frac{\phi-\gamma}{2}. \end{aligned} \tag{24}$$

Then one can change the η_a-integration into the γ_a-integration whose result is easily represented as the Kronecker delta δ_{m_1,m_2}. Hence, one can carry out m_2-summation and finally $(\vec{x}_b \mid \vec{x}_a)_E$ becomes

$$\begin{aligned} (\vec{x}_b \mid \vec{x}_a)_E = & \frac{-iM^2\omega}{\pi} \sum_{m=-\infty}^{\infty} e^{im(\phi_b-\phi_a)} \\ & \times \int_0^\infty \frac{dq}{\sinh^2 q} e^{-\frac{\xi}{\omega}q - 2M\omega(r_a+r_b)\coth q} \\ & \times I_{|m+\alpha|}\left(\frac{4M\omega}{\sinh q}\sqrt{r_a r_b}\cos\frac{\theta_a}{2}\cos\frac{\theta_b}{2}\right) I_{|m+\alpha|}\left(\frac{4M\omega}{\sinh q}\sqrt{r_a r_b}\sin\frac{\theta_a}{2}\sin\frac{\theta_b}{2}\right). \end{aligned} \tag{25}$$

In order to obtain the bound-state spectrum one has to check the pole of $(\vec{x}_b \mid \vec{x}_a)_E$ carfully. Following the appendix of Ref.[13], one can derive the following integral representation.

$$\begin{aligned} & I_{|m+\alpha|}\left(\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\cos\frac{\theta_a}{2}\cos\frac{\theta_b}{2}\right) I_{|m+\alpha|}\left(\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\sin\frac{\theta_a}{2}\sin\frac{\theta_b}{2}\right) \\ & = e^{-i\frac{\pi}{2}|m+\alpha|} 2^{-3|m+\alpha|} \pi^{-\frac{1}{2}} \left[\Gamma(\frac{1}{2}+|m+\alpha|)\right]^{-1} \left(\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\sin\theta_a \sin\theta_b\right)^{|m+\alpha|} \\ & \times \int_0^\pi d\phi (\sin\phi)^{2|m+\alpha|} \left[\cos\frac{\delta}{2}\right]^{-|m+\alpha|} J_{|m+\alpha|}\left(i\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\cos\frac{\delta}{2}\right) \end{aligned} \tag{26}$$

where $\cos\delta \equiv \cos\theta_a \cos\theta_b + \sin\theta_a \sin\theta_b \cos\phi$. If one expands $J_{|m+\alpha|}$ in Eq.(26) as[13]

$$\begin{aligned}
&J_{|m+\alpha|}\left(\frac{4iM\omega\sqrt{r_a r_b}}{\sinh q}\cos\frac{\delta}{2}\right) \\
&= \sqrt{2}\pi \frac{\Gamma(1+2|m+\alpha|)}{\Gamma(1+|m+\alpha|)}\left(\cos\frac{\delta}{2}\right)^{|m+\alpha|}\left(\frac{4iM\omega\sqrt{r_a r_b}}{\sinh q}\right)^{-|m+\alpha|-1} \\
&\times \sum_{n=0}^{\infty}\sum_{l=0}^{n}\left[\frac{(1+2|m+\alpha|+2n)(2|m+\alpha|+2l)(n-l)!}{\Gamma(\frac{1}{2}+|m+\alpha|)\Gamma(1+2|m+\alpha|+n+l)}\right. \\
&\times J_{1+2|m+\alpha|+2n}\left(\frac{4iM\omega\sqrt{r_a r_b}}{\sinh q}\right)(\sin\theta_a \sin\theta_b)^{-|m+\alpha|}(\sin\phi)^{\frac{1}{2}-|m+\alpha|} \\
&\times \left. P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_a) P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_b) P_{n+|m+\alpha|-\frac{1}{2}}^{|m+\alpha|-\frac{1}{2}}(\cos\phi)\right],
\end{aligned} \tag{27}$$

where P_n^m is a usual associated Legendre function, the ϕ-integration in Eq.(26) is easily performed by using an integral formula[14]

$$\begin{aligned}
&\int_{-1}^{1} dx (1-x^2)^{\lambda-1} P_\nu^\mu(x) \\
&= \frac{\pi 2^\mu \Gamma(\lambda+\frac{\mu}{2})\Gamma(\lambda-\frac{\mu}{2})}{\Gamma(\lambda+\frac{\nu}{2}+\frac{1}{2})\Gamma(\lambda-\frac{\nu}{2})\Gamma(-\frac{\mu}{2}+\frac{\nu}{2}+1)\Gamma(-\frac{\mu}{2}-\frac{\nu}{2}+\frac{1}{2})}.
\end{aligned} \tag{28}$$

Hence, Eq.(26) is reduced to

$$\begin{aligned}
&I_{|m+\alpha|}\left(\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\cos\frac{\theta_a}{2}\cos\frac{\theta_b}{2}\right) I_{|m+\alpha|}\left(\frac{4M\omega\sqrt{r_a r_b}}{\sinh q}\sin\frac{\theta_a}{2}\sin\frac{\theta_b}{2}\right) \\
&= e^{-i\pi\left(|m+\alpha|+\frac{1}{2}\right)} \frac{\pi \sinh q}{2M\omega\sqrt{r_a r_b}} \\
&\times \sum_{n=0}^{\infty}\sum_{l=0}^{n}\left[\frac{\sin\frac{l\pi}{2}}{\frac{l\pi}{2}} \frac{(1+2|m+\alpha|+2n)(|m+\alpha|+l)(n-l)!}{\Gamma(1+2|m+\alpha|+n+l)\Gamma(1+|m+\alpha|+\frac{l}{2})\Gamma(1-|m+\alpha|-\frac{l}{2})}\right. \\
&\times \left. J_{1+2|m+\alpha|+2n}\left(\frac{4iM\omega\sqrt{r_a r_b}}{\sinh q}\right) P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_a) P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_b)\right].
\end{aligned} \tag{29}$$

By inserting Eq.(29) into Eq.(25) and after doing the q-integration the final form of the fixed-energy amplitude is

$$\begin{aligned}
(\vec{x}_b \mid \vec{x}_a)_E &= \frac{1}{8\omega r_a r_b}\sum_{m=-\infty}^{\infty}\sum_{n=0}^{\infty}\sum_{l=0}^{n}\left[e^{im(\phi_b-\phi_a)} e^{-i\pi(|m+\alpha|+\frac{1}{2})}\frac{2}{l\pi}\sin\frac{l\pi}{2}\right. \\
&\times \frac{(|m+\alpha|+l)(n-l)!\Gamma(1+|m+\alpha|+n+\frac{\xi}{2\omega})}{\Gamma(1+2|m+\alpha|+n+l)\Gamma(1+2|m+\alpha|+2n)\Gamma(1-|m+\alpha|-\frac{l}{2})\Gamma(1+|m+\alpha|+\frac{l}{2})} \\
&\times P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_a) P_{n+|m+\alpha|}^{l+|m+\alpha|}(\cos\theta_b) \\
&\times \left. W_{-\frac{\xi}{2\omega},\frac{1+2|m+\alpha|+2n}{2}}\left(4M\omega \mathrm{Max}(r_a,r_b)\right) M_{\frac{\xi}{2\omega},\frac{1+2|m+\alpha|+2n}{2}}\left(-4M\omega \mathrm{Min}(r_a,r_b)\right)\right],
\end{aligned} \tag{30}$$

where $W_{a,b}$ and $M_{a,b}$ are usual Whittaker functions.

The energy spectrum of ABC system is deduced from the poles of the Gamma function in numerator:

$$E_{n,n',m} = -\frac{M\xi^2}{2(1+|m+\alpha|+n+n')^2}, \quad (n, n' = 0, 1, 2, \cdots). \tag{31}$$

Our result on the energy spectrum agrees with those of Ref.[5] and Ref.[6].

In summary, we derived the exact propagator of ABC system by using pseudotime method and dimensional extension technique firstly used by Duru and Kleinert in Ref.[8]. Also the bound state energy spectrum is deduced. It might be straightforward to derive the exact propagator of two-dimensional ABC system if one use a Levi-Civita transformation which is a two-dimensional version of the KS transformation. If the propagator of two dimensional ABC system is derived, one can derive the propagator of spin-1/2 ABC system by incorporating the self-adjoint extension method into the path integral formalism which was suggested in Ref.[15]. The evaluation of spin-1/2 ABC propagator might be very helpful for the analysis of time-dependent anyon scattering and study on the statistical properties of anyon system. This work is in progress.

REFERENCES

[1] Y.Aharonov and D.Bohm, *Phys. Rev.* 115, 485 (1959).

[2] M.Peshkin and T.Tonomura, *The Aharonov-Bohm Effect* (Springer-Verlag, Berlin, 1989).

[3] F.Wilczek, *Phys. Rev. Lett.* 48, 1144 (1982).

[4] C.R.Hagen, *Phys. Rev.* D48, 5935 (1993); D.K.Park and Jae Geun Oh, *Phys. Rev.* D50, 7715 (1994); C.R.Hagen and D.K.Park, *Ann. Phys.* 251, 45 (1996).

[5] L.Chetouani, L.Guechi and T.F.Hammann, *J. Math. Phys.* 30, 665 (1989).

[6] M.Kibler and T.Negadi, *Phys. Lett.* A124, 42 (1987); Gh. E.Draganascu. C.Campigotto and M.Kibler, *Phys. Lett.* A170, 339 (1992).

[7] P.Kustaanheimo and E.Stiefel, *J. Reine Angew. Math.* 218, 204 (1965).

[8] I.H.Duru and H.Kleinert, *Phys. Lett.* B84, 185 (1979).

[9] A.Inomata, *Phys. Lett.* A87, 387 (1982); R.Ho and A.Inomata, *Phys. Rev. Lett.* 48, 231 (1982).

[10] H.Kleinert, *Path Integrals in Quantum Mechanics, Statistics and Polymer Physics*, (World Scientific, Singapore, 1995).

[11] A.Inomata, in *Path Summation: Achievements and Goals*, edited by S.Lundqvist, A.Ranfagni, V.Sa-yakanit, and L.S.Schulman (World Scientific, Singapore, 1987).

[12] H.Kleinert, quant-ph/9612040.

[13] M.Z.Carpio and A.Inomata, in *path integrals from meV to MeV*, edited by M.C.Gutzwiller, A.Inomata, J.R.Klauder and L.Streit (World Scientific, Singapore, 1986).

[14] I.S.Gradshteyn and I.M.Ryzhik, *Table of Integrals, Series and Products* (Academic

Press, NY, 1965).

[15] D.K.Park, *J. Math. Phys.* 36, 5453 (1995).